



3197	Cys*-Phe-Trp-(D)Trp-Lys-Thr-Phe-Gly(S2)-X
3205	Phe(C3)-Cys*-Phe-(D)Trp-Lys-Thr-Cys*-Phe(N3)-X
3207	(D)Phe-Cys*-Phe-Trp-(D)Trp-Lys-Thr-Phe-Gly(S2)-X
3229	Galactose-Dab*-Phe-Trp-(D)Trp-Lys-Thr-Phe-Gly(C3)-X

Another preferred embodiment has the general formula:



Y<sup>2</sup> is thioether, thioester or disulfide.

Another more preferred embodiment has the general formula:



X designates a terminal carboxy acid, amide or alcohol group;

113 (over)

R<sup>5</sup> is (D)- or (L)-Phe or (D)- or (L)-Ala;  
 R<sup>7</sup> is (D)- or (L)-Trp, (D)- or (L)-Phe, (D)- or (L)- 1Nal or (D)- or (L)- 2Nal, or  
 Tyr;  
 R<sup>10</sup> is Thr, Gly, Abu, Ser, Cys, Val, (D)- or (L)-Ala, or (D)- or (L)-Phe;  
 R<sup>12</sup> is Gly, Val, (D)- or (L)-Phe or is absent;  
 R<sup>13</sup> is (D)- or (L)-Phe or (D)- or (L)-Ala; and  
 Y<sup>2</sup> is amide, thioether, thioester or disulfide.

Page 26, please replace old Table 3 with the following new Table 3:

Table 3. The most preferred analogs.

PTR	Sequence
3171	Phe*-Phe-Phe-(D)Trp-(D)Lys-Phe(C2)-X
3113	Phe(C1)-Phe-Phe-(D)Trp-Lys-Phe(N2)-X
3123	Phe(C1)-Phe-Phe-(D)Trp-(D)Lys-Phe(N2)-X
3209	Phe(N2)-Tyr-(D)2Nal-Lys-Val-Gly(C2)-Thr-X
3183	Phe(N2)-Tyr-(D)Trp-Lys-Val-Gly(C2)-2Nal-X
3185	Phe(N2)-Tyr-(D)Trp-Lys-Val-Val-Gly(C2)-X
3201	Phe(N2)-Tyr-(D)Trp-Lys-Ser-2Nal-Gly(C2)-X
3203	Phe(N2)-Phe-(D)Trp-Lys-Thr-2Nal-Gly(C2)-X
3173	GABA*-Phe-Trp-(D)Trp-Lys-Thr-Phe-Gly(C3)-X
3197	Cys*-Phe-Trp-(D)Trp-Lys-Thr-Phe-Gly(S2)-X
3205	Phe(C3)-Cys*-Phe-(D)Trp-Lys-Thr-Cys*-Phe(N3)-X
3207	(D)Phe-Cys*-Phe-Trp-(D)Trp-Lys-Thr-Phe-Gly(S2)-X
3229	Galactose-Dab*-Phe-Trp-(D)Trp-Lys-Thr-Phe-Gly(C3)-X

IN THE CLAIMS:

A marked version of the claims showing the amendments is attached hereto as Appendix B.

Please amend Claims 13, 14, and 15 to read as follows:

13. The backbone cyclized somatostatin analog of claim 1 having the general formula
- 15:

115